



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

nial root system probably does not supply by capillarity the constant stream of water as does the capillarity of the soil. But I do not think it unreasonable to suppose that there is a degree of root activity which furnishes the necessary water. The cold being superficial the water in the surface of the cambium crystallizes, the dead periderm cracks, and through the rift the nascent laminate crystal pushes its way.

A specific variation in the root activity of different plants as related to different temperatures explains, I think, why *Cunila Mariana* of all the plants Mr. Ward mentions forms the crystals. I discovered also one other plant which produced these frost freaks, but as the subject was losing what had seemed to me at first its very serious aspect, I did not take the trouble to accurately determine either the species or genus of this additional frost weed. From the observations which I made at the time I can safely say that it was either some species of *Eupatorium* or *Vernonia*, more likely the latter. I regret now that I did not accurately determine the species.—GEO. F. ATKINSON, *Botanical Department, Cornell University*.

A hybrid Baptisia.—Several specimens of a *Baptisia* have been collected in the vicinity of Manhattan which can not be referred to any of the species of the genus. The two species occurring here are *B. australis*, characterized by its glabrous foliage and erect raceme of blue flowers, and *B. leucophæa*, with hairy foliage and a reclining raceme of cream-colored flowers. The specimens referred to are intermediate in all these characters, even to the party-colored flowers, and are apparently hybrids between the two species. Fruiting specimens have not been observed.—A. S. HITCHCOCK, *Agricultural College, Manhattan, Kansas*.